

Available online at www.sciencedirect.com

Biochimica et Biophysica Acta xx (2007) xxx–xxx

www.elsevier.com/locate/bbamem

Review

The redox chemistry of the Alzheimer's disease amyloid β peptideDanielle G. Smith^{a,c,d}, Roberto Cappai^{a,b,c,d}, Kevin J. Barnham^{a,c,d,*}^a Department of Pathology, The University of Melbourne, Parkville, Victoria 3010, Australia^b Centre for Neuroscience, The University of Melbourne, Parkville, Victoria 3010, Australia^c Bio21 Institute, The University of Melbourne, Parkville, Victoria 3010, Australia^d The Mental Health Research Institute of Victoria, Parkville, Victoria 3052, Australia

Received 3 October 2006; received in revised form 2 February 2007; accepted 5 February 2007

Abstract

There is a growing body of evidence to support a role for oxidative stress in Alzheimer's disease (AD), with increased levels of lipid peroxidation, DNA and protein oxidation products (HNE, 8-HO-guanidine and protein carbonyls respectively) in AD brains. The brain is a highly oxidative organ consuming 20% of the body's oxygen despite accounting for only 2% of the total body weight. With normal ageing the brain accumulates metals ions such iron (Fe), zinc (Zn) and copper (Cu). Consequently the brain is abundant in antioxidants to control and prevent the detrimental formation of reactive oxygen species (ROS) generated via Fenton chemistry involving redox active metal ion reduction and activation of molecular oxygen. In AD there is an over accumulation of the Amyloid β peptide ($A\beta$), this is the result of either an elevated generation from amyloid precursor protein (APP) or inefficient clearance of $A\beta$ from the brain. $A\beta$ can efficiently generate reactive oxygen species in the presence of the transition metals copper and iron in vitro. Under oxidative conditions $A\beta$ will form stable dityrosine cross-linked dimers which are generated from free radical attack on the tyrosine residue at position 10. There are elevated levels of urea and SDS resistant stable linked $A\beta$ oligomers as well as dityrosine cross-linked peptides and proteins in AD brain. Since soluble $A\beta$ levels correlate best with the degree of degeneration [C.A. McLean, R.A. Cherny, F.W. Fraser, S.J. Fuller, M.J. Smith, K. Beyreuther, A.I. Bush, C.L. Masters, Soluble pool of Abeta amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease, *Ann. Neurol.* 46 (1999) 860–866] we suggest that the toxic $A\beta$ species corresponds to a soluble dityrosine cross-linked oligomer. Current therapeutic strategies using metal chelators such as clioquinol and desferrioxamine have had some success in altering the progression of AD symptoms. Similarly, natural antioxidants curcumin and ginkgo extract have modest but positive effects in slowing AD development. Therefore, drugs that target the oxidative pathways in AD could have genuine therapeutic efficacy.

© 2007 Elsevier B.V. All rights reserved.

Keywords: Amyloid beta; Metal; Oxidative stress; Redox chemistry; Alzheimer's disease**Contents**

1. Amyloid beta peptide ($A\beta$) generation	0
2. Oxidative stress in AD	0
3. Metals in AD	0
4. $A\beta$ is a metalloprotein	0
5. $A\beta$, metals and redox chemistry	0
5.1. The role of methionine 35 in $A\beta$ toxicity	0
5.2. A model for $A\beta$ redox chemistry	0
5.3. Oxidatively modified stable oligomeric $A\beta$ products	0

* Corresponding author. Department of Pathology, The University of Melbourne, Parkville, Victoria 3010, Australia. Tel.: +61 3 8344 2555; fax: +61 3 9347 6750.
E-mail address: kbarnham@unimelb.edu.au (K.J. Barnham).

6. Therapeutic inventions.	0
6.1. Antioxidants	0
6.2. Metal coordinating compounds.	0
7. Conclusions	0
Acknowledgements	0
References	0

1. Amyloid beta peptide (A β) generation

The A β peptide was first purified and sequenced from amyloid plaques found in AD and Down's syndrome brain [2,3]. It is typically a 39–42 residue polypeptide derived from the proteolytic processing of the APP molecule and consists of a largely hydrophilic N-terminal domain (1–28) and a C-terminal hydrophobic domain (29–39/43) derived from the APP transmembrane domain (Fig. 1). APP can be processed via two proteolytic pathways, each resulting in distinct cleavage products. Cleavage of APP by β -secretase generates a membrane bound C-terminal fragment which is the substrate for γ -secretase, which cleaves within the transmembrane domain of the C-terminal APP fragment to generate full length A β (Fig. 1). The alternative pathway involves initial cleavage of APP by α -secretase, followed by γ -secretase resulting in the truncated P3 fragment (Fig. 1).

APP is a ubiquitously expressed type 1 transmembrane protein and is a member of a multigene family which includes amyloid precursor-like proteins 1 and 2 (APLP1 and APLP2) [4–8]. The biological function of the APP family remains unknown. APP exhibits a variety of potential biological functions including regulation of intracellular calcium [9,10], cell growth [11–13], cell adhesion [14–17], axonal transport of vesicles [18] and metal ion homeostasis [19–21]. APP

contains a metal binding domain in the N-terminal cysteine-rich region adjacent to the growth factor-like domain [11,19–21]. Structural studies indicate the metal binding domain possesses structural similarities to known copper chaperones such as the Menkes copper-transporting ATPase fragment, metallochaperone Atx1, and SOD1 copper chaperone [21]. The APP copper binding domain exhibits a strong affinity for Cu²⁺ ions ($K_d \approx 10$ nM) and can reduce Cu²⁺ ions in vitro [19,20,22]. APP and APLP2 knockout mice have elevated Cu levels in the brain cortex and liver [23] while APP overexpressing mice exhibit a reduction in metals including copper ions [24–26].

2. Oxidative stress in AD

The brain is the most aerobically active organ in the body due its high metabolic requirements. The brain accounts for 2% of total body mass yet consumes 20% of total oxygen in a resting individual. Therefore it is imperative to maintain oxidative balance and control in the brain, and this is tightly regulated by antioxidants that are present in vastly higher amounts than in any other organ. Therefore modifications in normal oxidative metabolism as observed in AD brain provide strong evidence that oxidative stress plays an important role in AD pathogenesis (Fig. 2).




Fig. 1. Amyloid beta peptide (A β), a 39–43 amino acid transmembrane peptide is generated from APP cleavage by β - and γ -secretase. α -secretase cleavage of APP initiates the non-amyloidogenic processing pathway of APP. APP in a multi-domain protein consisting of eight domains: heparin binding (heparin), metal binding (metal), acidic, Kunitz-type serine protease inhibitor (KPI), OX-2 homology sequence (OX-2), carbohydrate, transmembrane (TM) and the cytoplasmic (cyto) domains [217]. The cleavage sites of the secretases are indicated by the arrows above the A β sequence. The transmembrane domain of A β is highlighted in orange and the familial mutations of A β are indicated below the sequence.




Fig. 2. A β reduction of Cu²⁺ ions generates A β radicals (A β •) that extract protons from surrounding lipids and proteins generating hydroxy-2-nonenal (HNE) and carbonyls respectively. Cu⁺ reacts with molecular oxygen (O₂) eventuating in H₂O₂ formation.

As a general principle, the chemical origin of the majority of ROS is the reaction of molecular oxygen with the redox active metals Cu and Fe [27]. The ability of these metal ions to occupy multiple valence states and undertake facile redox cycling, thereby activating molecular oxygen, has been utilized by a variety of enzymes including ceruloplasmin [28], cytochrome *c* oxidase [28] and amine oxidases [29]. However, unregulated redox-active metals will inappropriately react with oxygen to generate ROS. Moreover, there is increased glucose-6-phosphate dehydrogenase activity [30,31], increased heme oxygenase levels in diseased brain [32] and the presence of oxygen radical-mediated chemical attack resulting in increased free carbonyls, lipid peroxidation adducts, protein nitration and mitochondrial and nuclear DNA oxidation adducts.

A broad range of lipid peroxidation products can evolve from the cascade of events described above. High levels of chemically reactive electrophilic aldehydes malondialdehyde and free 4-hydroxy-2-nonenal (HNE) have been found in AD brain tissue [33–35]. These aldehydes readily react with cellular nucleophiles such as DNA, proteins and other lipids and elevated HNE-protein adducts have been detected in human AD brain [34,36,37]. They are potentially more perilous than free radicals because of their longer half-lives which allow them to diffuse away from their site of formation [38]. Polyunsaturated fatty acids, such as arachidonic acid and docosahexanoic acid, found in high abundance in brain [39] are highly oxidisable and hence vulnerable constituents of membrane phospholipids. Taken together, the above studies suggest that in an AD brain free radicals could induce lipid peroxidation leading to highly reactive aldehydes that initiate a cascade of oxidation events resulting in cellular dysfunction and ultimately death. There is significant evidence to indicate that A β can generate these free radicals (see later section).

Protein carbonyls are a major marker of protein oxidation and can be generated from direct free radical attack on amino acid side chains, glycation, glycooxidation or from lipid oxidation products as discussed above. Elevated levels of protein carbonyl exist in the frontal pole, hippocampus and superior middle temporal gyrus of AD patients and correlate well with AD histopathology [40–45]. Nitrotyrosine and dityrosine cross-linked protein are elevated 8-fold and 3-fold

respectively in hippocampus and neocortical regions of AD brain compared to age matched controls [26,46,47]. Both of these oxidation products in mammals generally correlate with oxidative stress [48–50]. Furthermore, neuritic and cored amyloid plaques show evidence of oxidatively modified A β [51] with much of the A β isolated from plaque containing methionine sulfoxide [52].

Further evidence of oxidative stress in AD is the modification of antioxidant activity in the brain. Thioredoxin is an antioxidant protein which is decreased in AD amygdala and hippocampus/parahippocampal gyrus. In contrast, the thioredoxin protein reductase shows increased activity in the amygdala and cerebellum of AD patients [53]. The antioxidant enzyme glutathione transferase, which is primarily responsible for HNE clearance, is decreased in several regions in AD brain including the hippocampus [54]. Furthermore, conjugated glutathione and HNE are elevated in the substantia innominata and the hippocampus in severe AD cases [55]. These two regions of the brain are known to be preferentially affected by the disease. HNE conjugation to glutathione by glutathione transferase is a cellular mechanism to inactivate toxic HNE. Therefore, increased levels would indicate an elevated cellular defence against lipid peroxidation.

The antioxidant enzyme superoxide dismutase (SOD) displays elevated expression levels in AD brains compared to age matched controls [56]. It is also elevated in red blood cells and lymphoblasts [57–60] which suggests that the normal compensatory mechanisms against free radical damage may be insufficient in AD patients. Apolipoprotein allele ϵ 4 increases the risk of developing AD and it has been shown that AD patients homozygous for this allele have an increase in brain catalase activity [61] suggesting a genetic predisposition for increased ROS generation in some AD suffers. Furthermore, the administration of the antioxidant vitamins ascorbic acid (vitamin C) and α -tocopherol (vitamin E) in tandem caused a decrease in the risk of AD in elderly subjects [62]. Taken together these observations suggest that prophylactic intake of antioxidants may be beneficial for those with a genetic risk of AD.

3. Metals in AD

The brain tightly regulates metal ion homeostasis as a part of normal physiological processes that play an important role in neural activity. Various lines of evidence have implicated metal ions, in particular Cu, Zn and Fe in the pathogenesis of AD (reviewed in [63]).

Transition metal ions Cu and Zn co-purify with A β extracted from AD plaques [64]. Cu ions are normally bound to Cu enzymes or proteins like cytochrome *c* oxidase, ceruloplasmin and superoxide dismutase during normal physiological circulation and are released into the synapse upon presynaptic excitation reaching up to 15 μ M in the synaptic cleft [65]. Although found in high concentrations in amyloid plaques (~400 μ M) [66] compared to the normal brain extracellular concentration of 0.2–1.7 μ M [67–72], most studies of bulk tissue have found either no statistical difference [73,74] or a

decrease [75,76] in total Cu concentration in AD brain compared to age matched controls.

Similarly, Zn ion levels can be as high as ~1 mM in plaques [66]. Interestingly, only mature cored senile plaques, not diffuse plaques, showed histochemical reactivity for Zn [77]. Using inductively coupled plasma mass spectrometry (ICP-MS), increased Zn levels were detected in cerebrospinal fluid (CSF) of AD patients [78]. There is conflicting data on brain Zn in AD levels with one report of bulk brain analysis indicating decreased levels of Zn in post mortem AD hippocampus [79], while a separate study observed an increase in the same region as well as in the amygdala and inferior parietal lobule [76,80]. Histological staining of normal brain indicates the distribution of Zn resembles the areas of the brain most prone to amyloid deposition and neuropathy in AD which includes the aforementioned hippocampus, amygdala and parietal lobe [81]. The primary source of labile Zn in the brain is from Zn released into the synapse during transmission [82] at ~200–300 μM [83,84]. Zn ions can promote A β aggregation and plaque formation, and this activity may not be as a neurotoxic modulator but rather as a neuroprotective agent since Zn can attenuate A β toxicity in cortical cultures [85–89]. The precise mechanism of cytoprotection is not clear, though possible mechanisms include inherent blockading of membrane channels formed [90–96], Zn enhancement of Na⁺/K⁺-ATPase [86] or competing with Cu for A β binding and thereby inhibiting A β initiated redox chemistry [85,97,98]. Oral Zn supplements significantly delay cognitive decline in AD patients [99–101].

The third transition metal found localized in human amyloid plaques is Fe. Bulk analysis of post mortem AD brain indicates that Fe is predominantly localized in neocortical grey matter [102], amygdala [103] and olfactory tract [73]. Despite having a high concentration in AD plaques (~1 μM) [66], Fe ions are not likely to interact directly with A β in vivo. Although in vitro studies indicate that Fe is able to interact with A β [104–106], unlike the other two transition metals Fe does not co-purify with A β extracted from plaque [64] and is predominantly located in neuritic processes within the plaque itself associated with ferritin [107]. A number of studies suggest that Fe homeostasis is altered in AD [66,107–111], however this is likely to be a secondary effect via another process such as increased heme oxygenase activity in response to cellular oxidative stress [32] or a decrease in functional heme from A β binding to heme increasing free Fe levels [112].

APP knockout mice have an increase in brain Cu levels [23]. In contrast, the Tg2576 APP transgenic mouse model which overexpresses the human APP Swedish mutation and forms amyloid plaques [113], display significantly reduced brain Cu levels prior to the appearance of amyloid neuropathology [24]. Two other APP transgenic mouse models, the APP23 and TgCRND8 mice also had reduced Cu levels. Zn [77] and Fe [114] levels were also dysregulated in the Tg2576 mice with the Tg2576 plaques containing high levels of both ions. The formation of amyloid plaques is associated with Zn-mediated A β aggregation since crossing the Tg2576 mice with mice where the synaptic Zinc transport 3 protein was knocked out

reduced the cerebral amyloid burden by 50% compared to Tg2576 control mice [115].

Oxidative stress markers have been detected in 12 month old senescence accelerated prone (SAMP8) mice which accumulate brain A β and display learning and memory impairments, and have elevated levels of oxidative stress markers compared to 4 month old mice [116]. Treating the aged SAMP8 mice with antisense oligonucleotide targeted to the A β sequence in APP showed decreased levels of oxidative stress markers including protein carbonyls, 3-nitrotyrosine adducts and HNE in the brain [117]. These results support a link between A β and oxidative stress.

4. A β is a metalloprotein

Multivalent metal ions are fundamental to redox chemistry as they facilitate electron transfers during the redox process. A β is a metalloprotein that displays high affinity binding of Cu²⁺, Zn²⁺ and Fe³⁺ ions [118]. The metal binding site was initially mapped to the region between positions 6 and 28. Solution state NMR implicated three histidine residues (His₆, His₁₃ and His₁₄) in binding Cu²⁺. This is supported by EPR data which suggests a square planar configuration in a 3N1O co-ordination sphere [119]. The nitrogen atoms involved in this co-ordination are from the imidazole rings of the three histidines (Fig. 3). The identity of the oxygen is unclear and it may be the hydroxyl from Tyr₁₀. Other options include the oxygen atoms of the sidechain carboxylate of Glu₅ [120], N-terminal aspartate [120], the amino terminus itself [120,121] or possibly a water molecule [122].

A β possesses a strong positive formal reduction potential and rapidly reduces Cu²⁺ and Fe³⁺ to Cu⁺ and Fe²⁺, respectively [98]. Molecular oxygen is then trapped generating free radical and peroxide species via Fenton chemistry with the A β 42 species exhibiting the greatest activity [98] (Fig. 2). The chemistry of these reactions is discussed in more detail later in this review. The free radicals generated are suggested to be involved in the oxidation of the methionine 35 residue (Met₃₅)








Fig. 3. Model of Cu²⁺ coordination with A β in a 3N1O coordination sphere via His₆, His₁₃, His₁₄ through the imidazole nitrogens. The fourth oxygen donor ligand putatively is the hydroxyl of Tyr₁₀, side chain carboxylate or a backbone carbonyl.

[123–129]. The replacement of Met₃₅ with norleucine (sulphur of methionine is replaced with a methylene group) abolishes the oxidative chemistry and neurotoxic activity as compared to the wildtype peptide [130]. The free radicals could possibly initiate the oxidation of the sulfhydryl to a sulfuranyl radical cation, which can then abstract protons from other lipids (causing lipid peroxidation) or proteins (initiating protein oxidation).

5. Aβ, metals and redox chemistry



There is good evidence that metal ions mediate the oxidative stress mechanism of Aβ toxicity. Metal ions modulate Aβ toxicity in neuronal cultures, with Cu²⁺ ions increasing Aβ toxicity whilst Zn²⁺ ions attenuate it [85]. Likewise, Aβ dissolved in Fe³⁺ containing media is toxic to neurons whilst Aβ in Fe³⁺ free media is not toxic [131]. The difference in the activity of the metal ions was attributed to the redox capabilities of Cu²⁺ and Fe³⁺. Aβ/Cu(Fe) complexes are capable of generating reactive oxygen species such as H₂O₂ which mediates toxicity [98,132]. Synthetic Aβ incubated in the presence of the chelator desferrioxamine (Fe³⁺ chelator) exhibited decreased toxicity which was restored by the addition of 0.1 mM Fe³⁺ [104]. Synthetic Aβ contains significant concentrations of Cu and Fe both of which can mediate Aβ production of H₂O₂ [133]. The addition of the H₂O₂ scavenging enzyme catalase to the cell cultures inhibits Aβ toxicity and suggests that Aβ neurotoxicity is mediated by ROS generation [98,132,134–137]. A caveat is that catalase can bind to Aβ and thus exert a static effect independent of the H₂O₂ production [138]. Co-incubation of Aβ with metal chelators can also attenuates Aβ toxicity [134,139] (Fig. 4).

The production of H₂O₂ by Aβ is dependant on the presence of specific redox active metal ions [132,140]. The generated peroxide may degrade forming a highly reactive hydroxyl radical (ROS) via Fenton chemistry or a Haber–Weiss reaction outlined as outlined below [141,142].



Aβ generation of ROS requires reduction of metal ions (Cu or Fe) thus inducing oxidation of another moiety. Mass spectrometry has shown that Cu²⁺ ions are able to oxygenate

Aβ with the most likely candidate being the sulphur atom of methionine 35 (Met₃₅) [143,144]. It has been proposed that Met₃₅ serves as a source of electrons for the reduction of molecular oxygen to hydrogen peroxide [119,127,130,145–148] (Eqs. (1)–(5)).



consistent with the in vitro data, oxidised methionine Aβ (AβMet-ox) has been isolated from AD brain amyloid deposits [149,150] and furthermore is bound to Cu [52].

The redox properties of Aβ indicate it could function as both an antioxidant and a pro-oxidant under specific conditions. Aβ has been observed at low concentration (nM range) to have a neuroprotective and neurotrophic effect on neonatal cell cultures [151–156]. However, the condition(s) under which Aβ ceases to act as an anti-oxidant and function as a pro-oxidant are not clearly understood, although several lines of evidence indicate the activity is dependant on the concentration of the peptide. The concentrations of Aβ required to induce toxicity in vitro is in the micromolar range [89,134,139,157–162]. In cerebrospinal fluid where Aβ has been reported to act as an anti-oxidant, the concentration of the peptide is between 0.1 and 1 nM, while at higher concentrations this activity was ablated [163]. Similarly, anti-apoptotic activity was only observed at nanomolar levels and conversely at higher concentrations it was toxic [164]. There is a strong negative correlation between oxidative damage and Aβ deposition in AD brain [1,165]. The accumulation of Aβ is either from excessive production of Aβ or alternatively an impairment of Aβ clearance pathways. Zn displacement of Cu binding to Aβ could be a detoxification mechanism followed by deposition into plaques which may not be toxic. Neuronal cells exposed to Aβ42 in the presence of both Cu and Zn had lower toxicity than Aβ plus Cu, suggesting Zn suppressed the Cu-dependent formation of H₂O₂ and rescued the cells [85]. Therefore, localized excessive levels of Aβ and Cu may generate ROS concentrations exceeding the capacity of the normal oxidation defence system. Moreover, synaptosomes treated with Aβ results in the release of free fatty acids and this effect was inhibited by the antioxidant vitamin E [166].

5.1. The role of methionine 35 in Aβ toxicity

There is conflicting data regarding the role of Met₃₅ in Aβ neurotoxicity. Met₃₅ is located in the C-terminal portion of the Aβ peptide, within the putative transmembrane domain and is the most susceptible side chain to oxidation [167,168]. Investigation into the role of Met₃₅ in Aβ toxicity suggests



Fig. 4. Oxidation of methionine residue leads to the formation of methionine sulfoxide.

the methionine is paramount to the redox chemistry of the peptide [127]. The sulfur containing methionine could potentially donate electrons for metal ion reduction and thus initiate a redox chemistry cycle as discussed previously.

An amino acid substitution of Met₃₅ to valine increases A β toxicity compared to the WT peptide [139] whilst nor-isoleucine or cysteine amino acid substitution ablated it [169]. The oxidised methionine A β peptide (A β Met-ox) displayed no toxicity after a 24 h exposure [130], however longer incubations of A β Met-ox exerted similar toxicity to wildtype A β [134]. These results support the concept that A β mediated generation of ROS is initiated by the Met₃₅ residue. In order for Met₃₅ to facilitate ROS generation by reducing metal ions, Met₃₅ must come within ~ 19 Å distance of the acceptor atoms for the electron transfer to occur [170]. This might be achieved by either the folding of the N and C-terminal ends or via the fibrillization of the peptide resulting in the metal bound N-terminal end of one A β being stacked adjacent to the Met₃₅ in the C-terminal end of another A β . Under these conditions, Cu²⁺ induced radicalization of Met₃₅ could also induce the generation of a stable carbon centred radical on the peptide backbone (most likely glycine residues), which can subsequently participate in lipid peroxidation reactions [171]. This is consistent with elevated post mortem 4-hydroxynonenal (HNE) levels in AD brain [33,172].

An alternative mechanism to the aforementioned process for the generation of HNE is based on full length A β inserting into the lipid membrane via its hydrophobic C-terminus wherein the sulfuranyl radical abstracts a proton from an unsaturated bond in the membrane phospholipid generating a carbon centred radical on the lipid. The carbon centred radical can readily react with molecular oxygen (O₂) forming a peroxy free radical (HO[•]) which in turn initiates a cascade of events that amplify the original A β peptide free radical. The lipid peroxidation products, such as HNE, arise from the systematic breakdown of the lipids subsequent to the free radical attack [127,173,174].

The oxidation state of the peptide can also affect aggregation as synthetic A β Met-ox has a slower rate of fibrillization, presumably by disrupting the switch from small soluble oligomers to larger insoluble oligomers. By arresting the oligomerisation to dimers, A β is able to be cleared from the brain before any toxic effect is exerted [175]. Interestingly,

A β Met-ox is more soluble than wildtype A β and was unable to insert into membranes and cause pore formation. However, the A β Met-ox was still neurotoxic [134] indicating membrane insertion is not a pre-requisite for A β toxicity.

5.2. A model for A β redox chemistry

The origin of the electron(s) that cause a reduction in the oxidation state of the metal ion could be the peptide itself, from Met₃₅ or from biological reducing agents such as dopamine and ascorbate [64]. The presence of an external reductant would permit the catalytic cycling of Cu or Fe without any net oxidation of the peptide. We used density function theory (DFT) to propose a detailed reaction scheme of the catalytic cycle that generates H₂O₂ (Fig. 5) [135]. The model proposed that as a consequence of the formation of a reactive tyrosyl radical on Tyr₁₀, that dityrosine cross-linked A β would be generated giving rise to covalently linked soluble oligomers. The formation of tyrosyl radicals is known to induce lipid peroxidation, a feature well characterized in AD.

The DFT model was supported by experimental data for the catalytic production of H₂O₂ by A β /Cu²⁺ and was shown to resemble the catalytic activity of galactose oxidase (GO). The role of the tyrosine residue was tested by using A β with alanine substituted for Tyr₁₀ (A β Y10A) [135]. The A β Y10A readily formed oligomers but was not toxic to cortical cultures whereas the wildtype peptide containing dityrosine linked oligomers was neurotoxic. Moreover A β Y10A can catalytically produce H₂O₂ at half the rate observed for the WT peptide consistent the pivotal role of the tyrosine residue in the redox chemistry as predicted by the model [135].

Opazo and colleagues have addressed other aspects of the model experimentally, specifically the ability of biologically significant reductants to initiate the redox chemistry [64]. A β 42, coordinated with up to two equivalents of Cu²⁺, can generate H₂O₂ catalytically by utilizing biological reducing agents as substrates under conditions where the Cu²⁺ or reductant will not form H₂O₂ themselves. The redox activity observed was inhibited by the presence of anti-A β antibody, Zn²⁺ or chelators targeting Cu [64]. Moreover, the toxicity of A β to neuronal cell cultures was accentuated with the addition of one such reductant dopamine [64].




Fig. 5. A simplified model of the redox chemistry events of A β catalytic production of H₂O₂ with dityrosine A β generated as a side product to the reaction.

5.3. Oxidatively modified stable oligomeric A β products

In addition to A β methionine sulfoxide, a number of other adducts resulting from redox reactions can be generated. Synthetically oxidatively modified A β products include tyrosine modified with adducts such as DOPA, dopamine, dopamine quinine, dihydroxyindol and isodityrosine (shown in Fig. 6) [176]. A β extracted from AD plaques contains oxidative modifications such as oxidised Met₃₅ [51,149], modification of the histidines to 2-oxo-histidine [177] and oxidatively modified tyrosine adducts including 3,4-dihydroxyphenylalanine (DOPA), dopamine and dopamine quinine [46,177,178]. Tyrosine is particularly susceptible to free radical attack due the conjugated aromatic ring. Evidence of other dityrosine linked peptides/proteins and 3-nitrotyrosine (Fig. 6) within the neuronal lesions in AD brain has been reported [51]. In vitro A β 42 in the presence of Cu²⁺ and H₂O₂ forms dityrosine cross-linked oligomers [179]. In a separate study, the formation of dityrosine linkage in A β facilitated further peptide aggregation, leading to the formation of higher order oligomers [143].

The formation of dityrosine adducts in vivo is a generalized sign of oxidative stress and is a result of free radical reactions [180,181]. The formation of dityrosine arises from the reaction of two tyrosine residues, which form carbon centered radicals in the aromatic ring (Fig. 7). The formation of dityrosine results in the creation of a very stable, irreversible covalent bond [182]. Stable A β oligomers resistant to SDS, urea and formic acid treatment have been extracted from AD brain tissue [3,183,184] suggesting that the A β peptides are covalently linked. Furthermore, naturally occurring stable A β oligomers but not monomers or fibrils inhibit hippocampal LTP [185]. We hypothesize that the A β toxic species requires the oxidative modification of A β resulting from redox chemistry that leads to

the formation of covalently linked (dityrosine crosslinked) soluble A β oligomers.

6. Therapeutic inventions

6.1. Antioxidants

The use of antioxidants has proven to be a promising approach for slowing progression of AD by inhibiting oxidative stress damage linked to cognitive and functional decline. A large amount of literature exists in relation to the potential benefits of vitamin supplements as AD prophylactics. Extensive in vitro studies are being conducted and some appear promising, however tandem administration of α -tocopherol (Vitamin E) (Fig. 8a) and ascorbic acid (Vitamin C) (Fig. 8b) only had modest benefits in elderly subjects [62]. Gossypin (3,3',4',5,7,8-hexahydroxyflavone 8-glucoside) (Fig. 8c), a bioflavonoid from the yellow petals of hibiscus flowers, is a radical scavenger and can protect cortical cell cultures [186] from A β induced toxicity. Melatonin (Fig. 8d) is neuroprotective against A β toxicity in vitro [187] and displayed beneficial effects in experimental mouse models of AD, including improvement of cognitive function [188], anti-oxidative injury [188,189], anti-apoptosis [188], inhibition of β -amyloid (A β) deposition and A β fibril formation [190]. Ginkgo biloba-extract EGb 761 also exhibited neuroprotective effects in several mouse models [191] as well as maintaining and improving cognitive function in AD patients [192,193]. In contrast, a double-blind placebo controlled study found no effect on dementia in AD patients [194]. In vitro studies indicate that the ginkgo extract activity is due to inhibition of A β induced free radical generation and occurs in a dose dependant manner [195].

Curcumin (1,7-bis(4-hydroxy-3-methoxyphenyl)-1,6 heptadiene-2,5 dione) (Fig. 8e), is the polyphenolic yellow pigment




Fig. 6. Tyrosine adducts formed under oxidative conditions.




Fig. 7. Dityrosine formation under oxidative conditions.

in the commonly used turmeric spice in Indian curries and food preservation [196]. Interestingly the prevalence of AD between the ages of 70–79 years in India is 4.4-fold less than in the USA [197]. The compound is neuroprotective against A β toxicity in vitro [198] while also being anti-amyloidogenic [199,200]. Furthermore aged transgenic Tg2576 mice with high amyloid plaque load either fed or injected with curcumin had less brain amyloid load and plaque burden; and curcumin labelled plaques [201]. Spectrophotometric studies indicate that curcumin is able to bind the more readily redox reactive metals Cu and Fe but does not bind Zn and therefore acts as an antioxidant by chelating the redox active metal ions [202].

While the details of the mode of pathogenesis remains elusive antioxidants that act to soak up the dangerous free radicals probably have some protective value. However, since they do not address the underlying cause of radical generation (metal-induced redox chemistry) they may have limited value as therapeutics once the disease has started to progress. This is because the reactions that define oxidative stress are chain reactions and once initiated give rise to a growing cascade of

events and therefore it is unlikely that sufficiently high concentrations of drug can reach the active sites to effectively soak up the radicals being generated. Therefore, antioxidants may find their most beneficial role when used as prophylactics.

6.2. Metal coordinating compounds

An alternative approach is to target the initiating event in the generation of free radicals: that is to employ metal complexing agents to prevent the metal ions from participating in the redox chemistry of A β that leads to the oxidative stress in AD. An important property of a potential AD therapeutic is its ability to cross the blood brain barrier (BBB). This excludes a large number of common metal chelators as possibilities due to their hydrophilic nature.

Desferrioxamine (*N'*-[5-(acetyl-hydroxy-amino)pentyl]-*N*-[5-[3-(5-aminoopentyl-hydroxy-carbamoyl)propanoylamino]pentyl]-*N*-hydroxy-butane diamide) (Fig. 9b), is a specific Fe chelator with high affinity also for Cu, Zn and aluminium (Al) [203,204]. In one trial desferrioxamine slowed the progression




Fig. 8. The molecular structure of antioxidants that show potential as Alzheimer's disease therapeutic treatments. (a) α -tocopherol, (b) ascorbic acid, (c) gossypin, (d) melatonin and (e) curcumin.




Fig. 9. The molecular structure of metal binding compounds that show potential as Alzheimer's disease therapeutic treatments. (a) clioquinol, (b) desferrioxamine, (c) bicyclam analogue JKL169, (d) triene and (e) tricyclodecan-9-yl-xanthogenate.

of AD. As a part of this study, the Al levels were monitored. Although Al has received some interest due to high concentrations being found in amyloid plaques and intracellular neurofibrillary tangles of AD brain [205–208], no convincing evidence has been published to suggest a role of the metal in the development of the disease although it remains a controversial issue [209]. Unfortunately no other metals were examined so it is unclear as to whether the therapeutic effect of desferrioxamine was actually due to chelation of other metals such as Cu and Zn. Another drawback for desferrioxamine is that it is not orally bioavailable.

Derivatives of a 14-membered saturated tetramine have attracted some recent interest. Bicyclam analogue JKL169 (1,1'-xylyl bis-1,4,8,11 tetraaza cyclotetradecane) (Fig. 9c) was effective in reducing Cu levels in brain cortex and is able to maintain normal Cu levels in the blood, CSF and corpus callosum in rats [210]. A number of other chelators have been studied such as the lipophilic chelator DP109 that reduced the level of aggregated insoluble A β and conversely increased soluble forms [211]. The chelator triene (triethylenetetramine) (Fig. 9d), which is used to treat Wilson's disease (intracellular Cu accumulation disease) was ineffective in reducing cerebral plaque concentration. This is attributed to the compound's inability to cross the blood brain barrier [212]. Brain tissue pre-exposed to tricyclodecan-9-yl-xanthogenate (Fig. 9e) showed significant reduction in oxidative stress markers after exposure to A β [213] however further work is required to evaluate whether the compound has promise as potential therapeutic for AD treatment.

Oral treatment with the chelator clioquinol (CQ, 5-chloro-7-iodo-8-hydroxyquinoline) (a former antibiotic) (Fig. 9a) in

Tg2576 mice resulted in a reduction of cortical deposition of amyloid (49%) with an improvement in general health and weight parameters remaining stable compared to untreated mice [212]. This quinoline compound is able to cross BBB and actually increases the brain Cu and Zn levels in treated mice. The increase in metal ions is suggested to be a result of CQ-metal complexes forming in the intestinal tract and crossing the BBB [210]. Clioquinol was used as an antibiotic, but was removed from the market in 1971 as it was implicated in the development of subacute myelo-optico-neuropathy (SMON) in some patients in Japan. However the link between CQ administration and the development of SMON in these patients is circumstantial as 25% of the diagnosed patients did not receive CQ [214] and was likely due to a vitamin B-12 deficiency within that population exacerbated by CQ [212,215]. The positive animal data lead to CQ being trialled clinically to treated AD, the drug was administered at much lower concentrations than was used to treat bacterial infections [216]. CQ showed significant promise for the treatment of AD, with a small phase II clinical trial revealing positive effects of the drug administered orally in moderately severe AD patients with no signs of SMON [216].

7. Conclusions

The brain requires high metal ion concentrations to carry out its numerous functions. However the brain has a poor capacity to cope with oxidative stress. Aberrant interactions between A β and redox active metals could initiate a cascade of events resulting in oxidative stress and chemically modified forms of A β . These chemical modifications include the generation of soluble covalently-linked oligomers of A β that have been

linked to the toxicity of the peptide. A number of therapeutic strategies aimed at ameliorating or inhibiting A β induced redox chemistry, including the use of antioxidants and metal complexing agents are currently being investigated for their utility in treating AD. Therefore, defining the primary trigger for the oxidative stress in AD and the role of A β in this process should be critical to our understanding of AD pathogenesis.

Acknowledgements

This work was supported in part by the, National Health and Medical Research Council of Australia, Australian Research Council, and Prana Biotechnology Ltd.

References

- [1] C.A. McLean, R.A. Cherny, F.W. Fraser, S.J. Fuller, M.J. Smith, K. Beyreuther, A.I. Bush, C.L. Masters, Soluble pool of Abeta amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease, *Ann. Neurol.* 46 (1999) 860–866.
- [2] G.G. Glenner, C.W. Wong, Alzheimer's disease and Down's syndrome: sharing of a unique cerebrovascular amyloid fibril protein, *Biochem. Biophys. Res. Commun.* 122 (1984) 1131–1135.
- [3] C.L. Masters, G. Simms, N.A. Weinman, G. Multhaup, B.L. McDonald, K. Beyreuther, Amyloid plaque core protein in Alzheimer disease and Down syndrome, *Proc. Natl. Acad. Sci. U. S. A.* 82 (1985) 4245–4249.
- [4] J. Kang, H.G. Lemaire, A. Unterbeck, J.M. Salbaum, C.L. Masters, K.H. Grzeschik, G. Multhaup, K. Beyreuther, B. Muller-Hill, The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor, *Nature* 325 (1987) 733–736.
- [5] R.E. Tanzi, J.F. Gusella, P.C. Watkins, G.A. Bruns, P. St George-Hyslop, M.L. Van Keuren, D. Patterson, S. Pagan, D.M. Kurnit, R.L. Neve, Amyloid beta protein gene: cDNA, mRNA distribution, and genetic linkage near the Alzheimer locus, *Science* 235 (1987) 880–884.
- [6] D. Goldgaber, M.I. Lerman, O.W. McBride, U. Saffiotti, D.C. Gajdusek, Characterization and chromosomal localization of a cDNA encoding brain amyloid of Alzheimer's disease, *Science* 235 (1987) 877–880.
- [7] W. Wasco, K. Bupp, M. Magendantz, J.F. Gusella, R.E. Tanzi, F. Solomon, Identification of a mouse brain cDNA that encodes a protein related to the Alzheimer disease-associated amyloid beta protein precursor, *Proc. Natl. Acad. Sci. U. S. A.* 89 (1992) 10758–10762.
- [8] H.H. Slunt, G. Thinakaran, C. Von Koch, A.C. Lo, R.E. Tanzi, S.S. Sisodia, Expression of a ubiquitous, cross-reactive homologue of the mouse beta-amyloid precursor protein (APP), *J. Biol. Chem.* 269 (1994) 2637–2644.
- [9] M.P. Mattson, B. Cheng, A.R. Culwell, F.S. Esch, I. Lieberburg, R.E. Rydel, Evidence for excitoprotective and intraneuronal calcium-regulating roles for secreted forms of the beta-amyloid precursor protein, *Neuron* 10 (1993) 243–254.
- [10] E.A. Milward, R. Papadopoulos, S.J. Fuller, R.D. Moir, D. Small, K. Beyreuther, C.L. Masters, The amyloid protein precursor of Alzheimer's disease is a mediator of the effects of nerve growth factor on neurite outgrowth, *Neuron* 9 (1992) 129–137.
- [11] J. Rossjohn, R. Cappai, S.C. Feil, A. Henry, W.J. McKinstry, D. Galatis, L. Hesse, G. Multhaup, K. Beyreuther, C.L. Masters, M.W. Parker, Crystal structure of the N-terminal, growth factor-like domain of Alzheimer amyloid precursor protein, *Nat. Struct. Biol.* 6 (1999) 327–331.
- [12] T. Saitoh, M. Sundsmo, J.M. Roch, N. Kimura, G. Cole, D. Schubert, T. Oltersdorf, D.B. Schenk, Secreted form of amyloid beta protein precursor is involved in the growth regulation of fibroblasts, *Cell* 58 (1989) 615–622.
- [13] D.H. Small, V. Nurcombe, G. Reed, H. Clarris, R. Moir, K. Beyreuther, C.L. Masters, A heparin-binding domain in the amyloid protein precursor of Alzheimer's disease is involved in the regulation of neurite outgrowth, *J. Neurosci.* 14 (1994) 2117–2127.
- [14] E.J. Coulson, G.L. Barrett, E. Storey, P.F. Bartlett, K. Beyreuther, C.L. Masters, Down-regulation of the amyloid protein precursor of Alzheimer's disease by antisense oligonucleotides reduces neuronal adhesion to specific substrata, *Brain Res.* 770 (1997) 72–80.
- [15] E. Storey, K. Beyreuther, C.L. Masters, Alzheimer's disease amyloid precursor protein on the surface of cortical neurons in primary culture co-localizes with adhesion patch components, *Brain Res.* 735 (1996) 217–231.
- [16] E. Storey, T. Spurck, J. Pickett-Heaps, K. Beyreuther, C.L. Masters, The amyloid precursor protein of Alzheimer's disease is found on the surface of static but not activity motile portions of neurites, *Brain Res.* 735 (1996) 59–66.
- [17] K.C. Breen, M. Bruce, B.H. Anderton, Beta amyloid precursor protein mediates neuronal cell–cell and cell-surface adhesion, *J. Neurosci. Res.* 28 (1991) 90–100.
- [18] S. Gunawardena, L.S. Goldstein, Disruption of axonal transport and neuronal viability by amyloid precursor protein mutations in *Drosophila*, *Neuron* 32 (2001) 389–401.
- [19] L. Hesse, D. Beher, C.L. Masters, G. Multhaup, The beta A4 amyloid precursor protein binding to copper, *FEBS Lett.* 349 (1994) 109–116.
- [20] G. Multhaup, A. Schlicksupp, L. Hesse, D. Beher, T. Ruppert, C.L. Masters, K. Beyreuther, The amyloid precursor protein of Alzheimer's disease in the reduction of copper(II) to copper(I), *Science* 271 (1996) 1406–1409.
- [21] K.J. Barnham, W.J. McKinstry, G. Multhaup, D. Galatis, C.J. Morton, C.C. Curtain, N.A. Williamson, A.R. White, M.G. Hinds, R.S. Norton, K. Beyreuther, C.L. Masters, M.W. Parker, R. Cappai, Structure of the Alzheimer's disease amyloid precursor protein copper binding domain. A regulator of neuronal copper homeostasis, *J. Biol. Chem.* 278 (2003) 17401–17407.
- [22] G. Multhaup, T. Ruppert, A. Schlicksupp, L. Hesse, E. Bill, R. Pipkorn, C.L. Masters, K. Beyreuther, Copper-binding amyloid precursor protein undergoes a site-specific fragmentation in the reduction of hydrogen peroxide, *Biochemistry* 37 (1998) 7224–7230.
- [23] A.R. White, R. Reyes, J.F. Mercer, J. Camakaris, H. Zheng, A.I. Bush, G. Multhaup, K. Beyreuther, C.L. Masters, R. Cappai, Copper levels are increased in the cerebral cortex and liver of APP and APLP2 knockout mice, *Brain Res.* 842 (1999) 439–444.
- [24] C.J. Maynard, R. Cappai, I. Volitakis, R.A. Cherny, A.R. White, K. Beyreuther, C.L. Masters, A.I. Bush, Q.X. Li, Overexpression of Alzheimer's disease amyloid-beta opposes the age-dependent elevations of brain copper and iron, *J. Biol. Chem.* 277 (2002) 44670–44676.
- [25] T.A. Bayer, S. Schafer, A. Simons, A. Kemmling, T. Kamer, R. Tepest, A. Eckert, K. Schussel, O. Eikenberg, C. Sturchler-Pierrat, D. Abramowski, M. Staufenbiel, G. Multhaup, Dietary Cu stabilizes brain superoxide dismutase 1 activity and reduces amyloid Abeta production in APP23 transgenic mice, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 14187–14192.
- [26] A.L. Phinney, B. Drisaldi, S.D. Schmidt, S. Lugowski, V. Coronado, Y. Liang, P. Horne, J. Yang, J. Sekoulidis, J. Coomaraswamy, M.A. Chishti, D.W. Cox, P.M. Mathews, R.A. Nixon, G.A. Carlson, P. St George-Hyslop, D. Westaway, In vivo reduction of amyloid-beta by a mutant copper transporter, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 14193–14198.
- [27] B. Halliwell, J.M. Gutterer, Free Radicals in Biology and Medicine, Oxford Univ. Press, Oxford, 1999.
- [28] M. Arredondo, M.T. Nunez, Iron and copper metabolism, *Mol. Aspects Med.* 26 (2005) 313–327.
- [29] G. Floris, A.F. Agro, Amine Oxidases, in: W.J. Lennarz, M.D. Lane (Eds.), *Encyclopedia of Biological Chemistry*, vol. 1, Elsevier Inc., 2004, pp. 85–89.
- [30] R.N. Martins, C.G. Harper, G.B. Stokes, C.L. Masters, Increased cerebral glucose-6-phosphate dehydrogenase activity in Alzheimer's disease may reflect oxidative stress, *J. Neurochem.* 46 (1986) 1042–1045.
- [31] L. Balazs, M. Leon, Evidence of an oxidative challenge in the Alzheimer's brain, *Neurochem. Res.* 19 (1994) 1131–1137.

- [32] H.M. Schipper, Heme oxygenase expression in human central nervous system disorders, *Free Radic. Biol. Med.* 37 (2004) 1995–2011.
- [33] W.R. Markesbery, M.A. Lovell, Four-hydroxynonenal, a product of lipid peroxidation, is increased in the brain in Alzheimer's disease, *Neurobiol. Aging* 19 (1998) 33–36.
- [34] L.M. Sayre, D.A. Zelasko, P.L. Harris, G. Perry, R.G. Salomon, M.A. Smith, 4-Hydroxynonenal-derived advanced lipid peroxidation end products are increased in Alzheimer's disease, *J. Neurochem.* 68 (1997) 2092–2097.
- [35] T.I. Williams, B.C. Lynn, W.R. Markesbery, M.A. Lovell, Increased levels of 4-hydroxynonenal and acrolein, neurotoxic markers of lipid peroxidation, in the brain in Mild Cognitive Impairment and early Alzheimer's disease, *Neurobiol. Aging* 27 (2006) 1094–1099.
- [36] K.S. Montine, P.J. Kim, S.J. Olson, W.R. Markesbery, T.J. Montine, 4-hydroxy-2-nonenal pyrrole adducts in human neurodegenerative disease, *J. Neuropathol. Exp. Neurol.* 56 (1997) 866–871.
- [37] Y. Ando, T. Brannstrom, K. Uchida, N. Nyhlin, B. Nasman, O. Suhr, T. Yamashita, T. Olsson, M. El Salhy, M. Uchino, M. Ando, Histochemical detection of 4-hydroxynonenal protein in Alzheimer amyloid, *J. Neurol. Sci.* 156 (1998) 172–176.
- [38] E.R. Stadtman, B.S. Berlett, Reactive oxygen-mediated protein oxidation in aging and disease, *Chem. Res. Toxicol.* 10 (1997) 485–494.
- [39] R.M. Nitsch, J.K. Blusztajn, A.G. Pittas, B.E. Slack, J.H. Growdon, R.J. Wurtman, Evidence for a membrane defect in Alzheimer disease brain, *Proc. Natl. Acad. Sci. U. S. A.* 89 (1992) 1671–1675.
- [40] R. Pamplona, E. Dalfó, V. Ayala, M.J. Bellmunt, J. Prat, I. Ferrer, M. Portero-Otin, Proteins in human brain cortex are modified by oxidation, glycoxidation, and lipoxidation. Effects of Alzheimer disease and identification of lipoxidation targets, *J. Biol. Chem.* 280 (2005) 21522–21530.
- [41] K. Hensley, N. Hall, R. Subramaniam, P. Cole, M. Harris, M. Aksenov, M. Aksenova, S.P. Gabbita, J.F. Wu, J.M. Carney, et al., Brain regional correspondence between Alzheimer's disease histopathology and biomarkers of protein oxidation, *J. Neurochem.* 65 (1995) 2146–2156.
- [42] D.A. Butterfield, S.M. Yatin, C.D. Link, In vitro and in vivo protein oxidation induced by Alzheimer's disease amyloid beta-peptide (1–42), *Ann. N. Y. Acad. Sci.* 893 (1999) 265–268.
- [43] M.V. Aksenova, M.Y. Aksenov, R.M. Payne, J.Q. Trojanowski, M.L. Schmidt, J.M. Carney, D.A. Butterfield, W.R. Markesbery, Oxidation of cytosolic proteins and expression of creatine kinase BB in frontal lobe in different neurodegenerative disorders, *Dement. Geriatr. Cogn. Disord.* 10 (1999) 158–165.
- [44] M.Y. Aksenov, M.V. Aksenova, D.A. Butterfield, J.W. Geddes, W.R. Markesbery, Protein oxidation in the brain in Alzheimer's disease, *Neuroscience* 103 (2001) 373–383.
- [45] L. Lyras, N.J. Cairns, A. Jenner, P. Jenner, B. Halliwell, An assessment of oxidative damage to proteins, lipids, and DNA in brain from patients with Alzheimer's disease, *J. Neurochem.* 68 (1997) 2061–2069.
- [46] K. Hensley, M.L. Maitz, Z. Yu, H. Sang, W.R. Markesbery, R.A. Floyd, Electrochemical analysis of protein nitrotyrosine and dityrosine in the Alzheimer brain indicates region-specific accumulation, *J. Neurosci.* 18 (1998) 8126–8132.
- [47] M.A. Smith, P.L. Richey Harris, L.M. Sayre, J.S. Beckman, G. Perry, Widespread peroxynitrite-mediated damage in Alzheimer's disease, *J. Neurosci.* 17 (1997) 2653–2657.
- [48] T.G. Huggins, M.W. Staton, D.G. Dyer, N.J. Detorie, M.D. Walla, J.W. Baynes, S.R. Thorpe, α -Tyrosine and dityrosine concentrations in oxidized proteins and lens proteins with age, *Ann. N. Y. Acad. Sci.* 663 (1992) 436–437.
- [49] C. Leeuwenburgh, P.A. Hansen, J.O. Holloszy, J.W. Heinecke, Hydroxyl radical generation during exercise increases mitochondrial protein oxidation and levels of urinary dityrosine, *Free Radic. Biol. Med.* 27 (1999) 186–192.
- [50] C. Leeuwenburgh, P.A. Hansen, J.O. Holloszy, J.W. Heinecke, Oxidized amino acids in the urine of aging rats: potential markers for assessing oxidative stress in vivo, *Am. J. Physiol.* 276 (1999) R128–R135.
- [51] E. Head, I.T. Lott, D.H. Cribbs, C.W. Cotman, T.T. Rohn, Beta-amyloid deposition and neurofibrillary tangle association with caspase activation in Down syndrome, *Neurosci. Lett.* 330 (2002) 99–103.
- [52] J. Dong, C.S. Atwood, V.E. Anderson, S.L. Siedlak, M.A. Smith, G. Perry, P.R. Carey, Metal binding and oxidation of amyloid-beta within isolated senile plaque cores: Raman microscopic evidence, *Biochemistry* 42 (2003) 2768–2773.
- [53] M.A. Lovell, C. Xie, S.P. Gabbita, W.R. Markesbery, Decreased thioredoxin and increased thioredoxin reductase levels in Alzheimer's disease brain, *Free Radic. Biol. Med.* 28 (2000) 418–427.
- [54] M.A. Lovell, C. Xie, W.R. Markesbery, Decreased glutathione transferase activity in brain and ventricular fluid in Alzheimer's disease, *Neurology* 51 (1998) 1562–1566.
- [55] W. Volkel, T. Sicilia, A. Pahler, W. Gsell, T. Tatschner, K. Jellinger, F. Leblhuber, P. Riederer, W.K. Lutz, M.E. Gotz, Increased brain levels of 4-hydroxy-2-nonenal glutathione conjugates in severe Alzheimer's disease, *Neurochem. Int.* 48 (2006) 679–686.
- [56] D.L. Marcus, J.A. Strafaci, M.L. Freedman, Differential neuronal expression of manganese superoxide dismutase in Alzheimer's disease, *Med. Sci. Monit.* 12 (2006) BR8–BR14.
- [57] J. Snaedal, J. Kristinsson, S. Gunnarsdottir, Á. Ólafsdóttir, M. Baldvinsson, T. Johannesson, Copper, ceruloplasmin and superoxide dismutase in patients with Alzheimer's disease. a case-control study, *Dement. Geriatr. Cogn. Disord.* 9 (1998) 239–242.
- [58] M.E. De Leo, S. Borrello, M. Passantino, B. Palazzotti, A. Mordente, A. Daniele, V. Filippini, T. Galeotti, C. Masullo, Oxidative stress and overexpression of manganese superoxide dismutase in patients with Alzheimer's disease, *Neurosci. Lett.* 250 (1998) 173–176.
- [59] F.P. Zemlan, O.J. Thienhaus, H.B. Bosmann, Superoxide dismutase activity in Alzheimer's disease: possible mechanism for paired helical filament formation, *Brain Res.* 476 (1989) 160–162.
- [60] R. Perrin, S. Briancon, C. Jeandel, Y. Artur, A. Minn, F. Penin, G. Siest, Blood activity of Cu/Zn superoxide dismutase, glutathione peroxidase and catalase in Alzheimer's disease: a case-control study, *Gerontology* 36 (1990) 306–313.
- [61] C. Ramassamy, D. Averill, U. Beffert, S. Bastianetto, L. Theroux, S. Lussier-Cacan, J.S. Cohn, Y. Christen, J. Davignon, R. Quirion, J. Poirier, Oxidative damage and protection by antioxidants in the frontal cortex of Alzheimer's disease is related to the apolipoprotein E genotype, *Free Radic. Biol. Med.* 27 (1999) 544–553.
- [62] P.P. Zandi, J.C. Anthony, A.S. Khachaturian, S.V. Stone, D. Gustafson, J.T. Tschanz, M.C. Norton, K.A. Welsh-Bohmer, J.C. Breitner, Reduced risk of Alzheimer disease in users of antioxidant vitamin supplements: the Cache County Study, *Arch. Neurol.* 61 (2004) 82–88.
- [63] A.I. Bush, The metallobiology of Alzheimer's disease, *Trends Neurosci.* 26 (2003) 207–214.
- [64] C. Opazo, X. Huang, R.A. Cherny, R.D. Moir, A.E. Roher, A.R. White, R. Cappai, C.L. Masters, R.E. Tanzi, N.C. Inestrosa, A.I. Bush, Metalloenzyme-like activity of Alzheimer's disease beta-amyloid. Cu-dependent catalytic conversion of dopamine, cholesterol, and biological reducing agents to neurotoxic H(2)O(2), *J. Biol. Chem.* 277 (2002) 40302–40308.
- [65] D.E. Hartter, A. Barnea, Brain tissue accumulates 67copper by two ligand-dependent saturable processes. A high affinity, low capacity and a low affinity, high capacity process, *J. Biol. Chem.* 263 (1988) 799–805.
- [66] M.A. Lovell, J.D. Robertson, W.J. Teesdale, J.L. Campbell, W.R. Markesbery, Copper, iron and zinc in Alzheimer's disease senile plaques, *J. Neurol. Sci.* 158 (1998) 47–52.
- [67] J.M. Gutteridge, Copper-phenanthroline-induced site-specific oxygen-radical damage to DNA. Detection of loosely bound trace copper in biological fluids, *Biochem. J.* 218 (1984) 983–985.
- [68] J. Kardos, I. Kovacs, F. Hajos, M. Kalman, M. Simonyi, Nerve endings from rat brain tissue release copper upon depolarization. A possible role in regulating neuronal excitability, *Neurosci. Lett.* 103 (1989) 139–144.
- [69] M.C. Linder, M. Hazegh-Azam, Copper biochemistry and molecular biology, *Am. J. Clin. Nutr.* 63 (1996) 797S–811S.
- [70] K. Schumann, H.G. Classen, H.H. Dieter, J. König, G. Multhaup, M. Rukgauer, K.H. Summer, J. Bernhardt, H.K. Biesalski, Hohenheim consensus workshop: copper, *Eur. J. Clin. Nutr.* 56 (2002) 469–483.

- [71] H.J. Stuerenburg, CSF copper concentrations, blood–brain barrier function, and ceruloplasmin synthesis during the treatment of Wilson's disease, *J. Neural Transm.* 107 (2000) 321–329.
- [72] A.R. White, K.J. Barnham, X. Huang, I. Voltakis, K. Beyreuther, C.L. Masters, R.A. Cherny, A.I. Bush, R. Cappai, Iron inhibits neurotoxicity induced by trace copper and biological reductants, *J. Biol. Inorg. Chem.* 9 (2004) 269–280.
- [73] L. Tandon, B.-F. Ni, X.X. Ding, W.D. Ehmann, E.J. Kassarskis, W.R. Markesbery, RNAA for arsenic, cadmium, copper and molybdenum in CNS tissues from subjects with age-related neurodegenerative disease, *J. Radioanal. Nucl. Chem.* 179 (1994) 331–339.
- [74] N.I. Ward, J.A. Mason, Neutron activation analysis techniques for identifying elemental status in Alzheimer's disease, *J. Radioanal. Nucl. Chem.* 113 (1987) 515–526.
- [75] L.-O. Plantin, U. Lysing-Tunell, K. Kristensson, Trace elements in the human central nervous system studied with neutron activation analysis, *Biol. Trace Elem. Res.* 1987 (1987) 69–75.
- [76] M.A. Deibel, W.D. Ehmann, W.R. Markesbery, Copper, iron, and zinc imbalances in severely degenerated brain regions in Alzheimer's disease: possible relation to oxidative stress, *J. Neurol. Sci.* 143 (1996) 137–142.
- [77] J.Y. Lee, I. Mook-Jung, J.Y. Koh, Histochemically reactive zinc in plaques of the Swedish mutant beta-amyloid precursor protein transgenic mice, *J. Neurosci.* 19 (1999) RC10.
- [78] C.O. Hershey, L.A. Hershey, A. Varnes, S.D. Vibhakar, P. Lavin, W.H. Strain, Cerebrospinal fluid trace element content in dementia: clinical, radiologic, and pathologic correlations, *Neurology* 33 (1983) 1350–1353.
- [79] F.M. Corrigan, G.P. Reynolds, N.I. Ward, Hippocampal tin, aluminum and zinc in Alzheimer's disease, *Biometals* 6 (1993) 149–154.
- [80] M. Diebel, W. Ehrmann, W.R. Markesbery, Copper, Iron and zinc imbalances in severely degenerated brain regions in Alzheimer's disease: possible relation to oxidative stress, *J. Neurol. Sci.* 143 (1996) 137–142.
- [81] A.I. Bush, Metal complexing agents as therapies for Alzheimer's disease, *Neurobiol. Aging* 23 (2002) 1031–1038.
- [82] C.J. Frederickson, Neurobiology of zinc and zinc-containing neurons, *Int. Rev. Neurobiol.* 31 (1989) 145–238.
- [83] S.Y. Assaf, S.H. Chung, Release of endogenous Zn²⁺ from brain tissue during activity, *Nature* 308 (1984) 734–736.
- [84] G.A. Howell, M.G. Welch, C.J. Frederickson, Stimulation-induced uptake and release of zinc in hippocampal slices, *Nature* 308 (1984) 736–738.
- [85] M.P. Cuajungco, L.E. Goldstein, A. Nunomura, M.A. Smith, J.T. Lim, C.S. Atwood, X. Huang, Y.W. Farrag, G. Perry, A.I. Bush, Evidence that the beta-amyloid plaques of Alzheimer's disease represent the redox-silencing and entombment of abeta by zinc, *J. Biol. Chem.* 275 (2000) 19439–19442.
- [86] M.A. Lovell, C. Xie, W.R. Markesbery, Protection against amyloid beta peptide toxicity by zinc, *Brain Res.* 823 (1999) 88–95.
- [87] K.S. Fuson, L.N. Boggs, P.C. May, Zinc promotes Abeta aggregation but attenuates Abeta neurotoxicity, *Neurobiol. Aging* 17 (1996) 431.
- [88] Y. Yoshiike, K. Tanemura, O. Murayama, T. Akagi, M. Murayama, S. Sato, X. Sun, N. Tanaka, A. Takashima, New insights on how metals disrupt amyloid beta-aggregation and their effects on amyloid-beta cytotoxicity, *J. Biol. Chem.* 276 (2001) 32293–32299.
- [89] S.M. Cardoso, A.C. Rego, C. Pereira, C.R. Oliveira, Protective effect of zinc on amyloid β -25–35 and 1–40? mediated toxicity, *Neurotox. Res.* 7 (2005) 273–282.
- [90] R. Bhatia, H. Lin, R. Lal, Fresh and globular amyloid beta protein (1–42) induces rapid cellular degeneration: evidence for AbetaP channel-mediated cellular toxicity, *FASEB J.* 14 (2000) 1233–1243.
- [91] Y. Hirakura, W.W. Yiu, A. Yamamoto, B.L. Kagan, Amyloid peptide channels: blockade by zinc and inhibition by Congo red (amyloid channel block), *Amyloid* 7 (2000) 194–199.
- [92] H. Lin, Y.J. Zhu, R. Lal, Amyloid beta protein (1–40) forms calcium-permeable, Zn²⁺-sensitive channel in reconstituted lipid vesicles, *Biochemistry* 38 (1999) 11189–11196.
- [93] S.K. Rhee, A.P. Quist, R. Lal, Amyloid beta protein-(1–42) forms calcium-permeable, Zn²⁺-sensitive channel, *J. Biol. Chem.* 273 (1998) 13379–13382.
- [94] Y.J. Zhu, H. Lin, R. Lal, Fresh and nonfibrillar amyloid beta protein (1–40) induces rapid cellular degeneration in aged human fibroblasts: evidence for AbetaP-channel-mediated cellular toxicity, *FASEB J.* 14 (2000) 1244–1254.
- [95] N. Arispe, H.B. Pollard, E. Rojas, Zn²⁺ interaction with Alzheimer amyloid beta protein calcium channels, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 1710–1715.
- [96] M. Kawahara, N. Arispe, Y. Kuroda, E. Rojas, Alzheimer's disease amyloid beta-protein forms Zn(2+)-sensitive, cation-selective channels across excised membrane patches from hypothalamic neurons, *Biophys. J.* 73 (1997) 67–75.
- [97] X. Huang, C.S. Atwood, R.D. Moir, M.A. Hartshorn, J.P. Vonsattel, R.E. Tanzi, A.I. Bush, Zinc-induced Alzheimer's Abeta1–40 aggregation is mediated by conformational factors, *J. Biol. Chem.* 272 (1997) 26464–26470.
- [98] X. Huang, M.P. Cuajungco, C.S. Atwood, M.A. Hartshorn, J.D. Tyndall, G.R. Hanson, K.C. Stokes, M. Leopold, G. Multhaup, L.E. Goldstein, R.C. Scarpa, A.J. Saunders, J. Lim, R.D. Moir, C. Glabe, E.F. Bowden, C.L. Masters, D.P. Fairlie, R.E. Tanzi, A.I. Bush, Cu(II) potentiation of Alzheimer abeta neurotoxicity. Correlation with cell-free hydrogen peroxide production and metal reduction, *J. Biol. Chem.* 274 (1999) 37111–37116.
- [99] R. Tudor, P.D. Zalewski, R.N. Ratnaik, Zinc in health and chronic disease, *J. Nutr. Health Aging* 9 (2005) 45–51.
- [100] F.C. Potocnik, S.J. van Rensburg, C. Park, J.J. Taljaard, R.A. Emsley, Zinc and platelet membrane microviscosity in Alzheimer's disease. The in vivo effect of zinc on platelet membranes and cognition, *S. Afr. Med. J.* 87 (1997) 1116–1119.
- [101] A.G. Van Rhijin, C.A. Prior, F.M. Corrigan, Dietary Supplementation with zinc sulphate, sodium selenite and fatty acids in early dementia of Alzheimer's type, *J. Nutr. Med.* 1 (1990) 259–266.
- [102] W.D. Ehmann, W.R. Markesbery, M. Alauddin, T.I.M. Hossain, E.H. Brubaker, Brain Trace elements in Alzheimer's disease, *Neurotoxicology* 7 (1986) 197–206.
- [103] C.M. Thompson, W.R. Markesbery, W.D. Ehmann, Y.X. Mao, D.E. Vance, Regional brain trace element studies in Alzheimer's disease, *Neurotoxicology* 9 (1988) 1–8.
- [104] C.A. Rottkamp, A.K. Raina, X. Zhu, E. Gaier, A.I. Bush, C.S. Atwood, M. Chevion, G. Perry, M.A. Smith, Redox-active iron mediates amyloid-beta toxicity, *Free Radic. Biol. Med.* 30 (2001) 447–450.
- [105] A. Khan, J.P. Dobson, C. Exley, Redox cycling of iron by Abeta42, *Free Radic. Biol. Med.* 40 (2006) 557–569.
- [106] W.P. Hu, G.L. Chang, S.J. Chen, Y.M. Kuo, Kinetic analysis of beta-amyloid peptide aggregation induced by metal ions based on surface plasmon resonance biosensing, *J. Neurosci. Methods* 154 (2006) 190–197.
- [107] I. Grundke-Iqbal, J. Fleming, Y.C. Tung, H. Lassmann, K. Iqbal, J.G. Joshi, Ferritin is a component of the neuritic (senile) plaque in Alzheimer dementia, *Acta Neuropathol. (Berl.)* 81 (1990) 105–110.
- [108] R.C. Switzer, T.L. Martin, S.K. Campbell, J.C. Parker, E.D. Caldwell, Iron and ferritin in the neuritic plaques of Alzheimer's disease, *Abstr. - Soc. Neurosci.* 100 (1986).
- [109] J.R. Connor, B.S. Snyder, J.L. Beard, R.E. Fine, E.J. Mufson, Regional distribution of iron and iron-regulatory proteins in the brain in aging and Alzheimer's disease, *J. Neurosci. Res.* 31 (1992) 327–335.
- [110] J.R. Connor, S.L. Menzies, S.M. St Martin, E.J. Mufson, A histochemical study of iron, transferrin, and ferritin in Alzheimer's diseased brains, *J. Neurosci. Res.* 31 (1992) 75–83.
- [111] J.R. Connor, S.L. Menzies, Cellular management of iron in the brain, *J. Neurol. Sci.* 134 (1995) 33–44 (Suppl.).
- [112] H. Atamna, W.H. Frey II, A role for heme in Alzheimer's disease: heme binds amyloid beta and has altered metabolism, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 11153–11158.
- [113] K. Hsiao, P. Chapman, S. Nilsen, C. Eckman, Y. Harigaya, S. Younkin, F. Yang, G. Cole, Correlative memory deficits, Abeta elevation, and amyloid plaques in transgenic mice, *Science* 274 (1996) 99–102.
- [114] M.A. Smith, K. Wehr, P.L. Harris, S.L. Siedlak, J.R. Connor, G. Perry,

- Abnormal localization of iron regulatory protein in Alzheimer's disease, *Brain Res.* 788 (1998) 232–236.
- [115] J.Y. Lee, T.B. Cole, R.D. Palmiter, S.W. Suh, J.Y. Koh, Contribution by synaptic zinc to the gender-disparate plaque formation in human Swedish mutant APP transgenic mice, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 7705–7710.
- [116] J.F. Flood, J.E. Morley, Age-related changes in footshock avoidance acquisition and retention in senescence accelerated mouse (SAM), *Neurobiol. Aging* 14 (1993) 153–157.
- [117] H.F. Poon, G. Joshi, R. Sultana, S.A. Farr, W.A. Banks, J.E. Morley, V. Calabrese, D.A. Butterfield, Antisense directed at the Abeta region of APP decreases brain oxidative markers in aged senescence accelerated mice, *Brain Res.* 1018 (2004) 86–96.
- [118] C.S. Atwood, R.C. Scarpa, X. Huang, R.D. Moir, W.D. Jones, D.P. Fairlie, R.E. Tanzi, A.I. Bush, Characterization of copper interactions with Alzheimer amyloid beta peptides: identification of an atomolar-affinity copper binding site on amyloid beta1–42, *J. Neurochem.* 75 (2000) 1219–1233.
- [119] C.C. Curtain, F. Ali, I. Volitakis, R.A. Cherny, R.S. Norton, K. Beyreuther, C.J. Barrow, C.L. Masters, A.I. Bush, K.J. Barnham, Alzheimer's disease amyloid-beta binds copper and zinc to generate an allosterically ordered membrane-penetrating structure containing superoxide dismutase-like subunits, *J. Biol. Chem.* 276 (2001) 20466–20473.
- [120] J.W. Karr, H. Akintoye, L.J. Kaupp, V.A. Szalai, N-terminal deletions modify the Cu²⁺ binding site in amyloid-beta, *Biochemistry* 44 (2005) 5478–5487.
- [121] C.D. Syme, R.C. Nadal, S.E. Rigby, J.H. Viles, Copper binding to the amyloid-beta (Abeta) peptide associated with Alzheimer's disease: folding, coordination geometry, pH dependence, stoichiometry, and affinity of Abeta(1–28): insights from a range of complementary spectroscopic techniques, *J. Biol. Chem.* 279 (2004) 18169–18177.
- [122] T. Miura, K. Suzuki, N. Kohata, H. Takeuchi, Metal binding modes of Alzheimer's amyloid beta-peptide in insoluble aggregates and soluble complexes, *Biochemistry* 39 (2000) 7024–7031.
- [123] D. Boyd-Kimball, R. Sultana, H. Mohmmad-Abdul, D.A. Butterfield, Rodent Abeta(1–42) exhibits oxidative stress properties similar to those of human Abeta(1–42): Implications for proposed mechanisms of toxicity, *J. Alzheimer's Dis.* 6 (2004) 515–525.
- [124] J. Kanski, M. Aksenova, C. Schoneich, D.A. Butterfield, Substitution of isoleucine-31 by helical-breaking proline abolishes oxidative stress and neurotoxic properties of Alzheimer's amyloid beta-peptide, *Free Radic. Biol. Med.* 32 (2002) 1205–1211.
- [125] D.A. Butterfield, J. Kanski, Methionine residue 35 is critical for the oxidative stress and neurotoxic properties of Alzheimer's amyloid beta-peptide 1–42, *Peptides* 23 (2002) 1299–1309.
- [126] S.M. Yatin, S. Varadarajan, C.D. Link, D.A. Butterfield, In vitro and in vivo oxidative stress associated with Alzheimer's amyloid beta-peptide (1–42), *Neurobiol. Aging* 20 (1999) 325–330 discussion 339–342.
- [127] D.A. Butterfield, Amyloid beta-peptide (1–42)-induced oxidative stress and neurotoxicity: implications for neurodegeneration in Alzheimer's disease brain. A review, *Free Radical Res.* 36 (2002) 1307–1313.
- [128] D.A. Butterfield, A.I. Bush, Alzheimer's amyloid beta-peptide (1–42): involvement of methionine residue 35 in the oxidative stress and neurotoxicity properties of this peptide, *Neurobiol. Aging* 25 (2004) 563–568.
- [129] D.A. Butterfield, S. Griffin, G. Munch, G.M. Pasinetti, Amyloid beta-peptide and amyloid pathology are central to the oxidative stress and inflammatory cascades under which Alzheimer's disease brain exists, *J. Alzheimers Dis.* 4 (2002) 193–201.
- [130] S. Varadarajan, J. Kanski, M. Aksenova, C. Lauderback, D.A. Butterfield, Different mechanisms of oxidative stress and neurotoxicity for Alzheimer's A beta(1–42) and A beta(25–35), *J. Am. Chem. Soc.* 123 (2001) 5625–5631.
- [131] D. Schubert, M. Chevion, The role of iron in beta amyloid toxicity, *Biochem. Biophys. Res. Commun.* 216 (1995) 702–707.
- [132] X. Huang, C.S. Atwood, M.A. Hartshorn, G. Multhaup, L.E. Goldstein, R.C. Scarpa, M.P. Cuajungco, D.N. Gray, J. Lim, R.D. Moir, R.E. Tanzi, A.I. Bush, The A beta peptide of Alzheimer's disease directly produces hydrogen peroxide through metal ion reduction, *Biochemistry* 38 (1999) 7609–7616.
- [133] S. Turnbull, B.J. Tabner, O.M. El-Agnaf, L.J. Twyman, D. Allsop, New evidence that the Alzheimer beta-amyloid peptide does not spontaneously form free radicals: an ESR study using a series of spin-traps, *Free Radic. Biol. Med.* 30 (2001) 1154–1162.
- [134] K.J. Barnham, G.D. Ciccotosto, A.K. Tickler, F.E. Ali, D.G. Smith, N.A. Williamson, Y.H. Lam, D. Carrington, D. Tew, G. Kocak, I. Volitakis, F. Separovic, C.J. Barrow, J.D. Wade, C.L. Masters, R.A. Cherny, C.C. Curtain, A.I. Bush, R. Cappai, Neurotoxic, redox-competent Alzheimer's beta-amyloid is released from lipid membrane by methionine oxidation, *J. Biol. Chem.* 278 (2003) 42959–42965.
- [135] K.J. Barnham, F. Haeffner, G.D. Ciccotosto, C.C. Curtain, D. Tew, C. Mavros, K. Beyreuther, D. Carrington, C.L. Masters, R.A. Cherny, R. Cappai, A.I. Bush, Tyrosine gated electron transfer is key to the toxic mechanism of Alzheimer's disease beta-amyloid, *FASEB J.* 18 (2004) 1427–1429.
- [136] A.J. Bruce, B. Malfroy, M. Baudry, beta-Amyloid toxicity in organotypic hippocampal cultures: protection by EUK-8, a synthetic catalytic free radical scavenger, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 2312–2316.
- [137] C. Behl, J.B. Davis, R. Lesley, D. Schubert, Hydrogen peroxide mediates amyloid beta protein toxicity, *Cell* 77 (1994) 817–827.
- [138] N.G. Milton, Amyloid-beta binds catalase with high affinity and inhibits hydrogen peroxide breakdown, *Biochem. J.* 344 (Pt 2) (1999) 293–296.
- [139] G.D. Ciccotosto, D. Tew, C.C. Curtain, D. Smith, D. Carrington, C.L. Masters, A.I. Bush, R.A. Cherny, R. Cappai, K.J. Barnham, Enhanced toxicity and cellular binding of a modified amyloid beta peptide with a methionine to valine substitution, *J. Biol. Chem.* 279 (2004) 42528–42534.
- [140] B.J. Tabner, S. Turnbull, O.M. El-Agnaf, D. Allsop, Formation of hydrogen peroxide and hydroxyl radicals from A(beta) and alpha-synuclein as a possible mechanism of cell death in Alzheimer's disease and Parkinson's disease, *Free Radic. Biol. Med.* 32 (2002) 1076–1083.
- [141] W.R. Markesbery, Oxidative stress hypothesis in Alzheimer's disease, *Free Radic. Biol. Med.* 23 (1997) 134–147.
- [142] S.I. Liochev, The mechanism of "Fenton-like" reactions and their importance for biological systems. A biologist's view, *Met. Ions Biol. Syst.* 36 (1999) 1–39.
- [143] F.E. Ali, F. Separovic, C.J. Barrow, R.A. Cherny, F. Fraser, A.I. Bush, C.L. Masters, K.J. Barnham, Methionine regulates copper/hydrogen peroxide oxidation products of Abeta, *J. Pept. Sci.* 11 (2005) 353–360.
- [144] A.J. Schiewe, L. Margol, B.A. Soreghan, S.N. Thomas, A.J. Yang, Rapid characterization of amyloid-beta side-chain oxidation by tandem mass spectrometry and the scoring algorithm for spectral analysis, *Pharm. Res.* 21 (2004) 1094–1102.
- [145] M.E. Clementi, S. Marini, M. Coletta, F. Orsini, B. Giardina, F. Misiti, Abeta(31–35) and Abeta(25–35) fragments of amyloid beta-protein induce cellular death through apoptotic signals: Role of the redox state of methionine-35, *FEBS Lett.* 579 (2005) 2913–2918.
- [146] F. Misiti, G.E. Martorana, G. Nocca, E. Di Stasio, B. Giardina, M.E. Clementi, Methionine 35 oxidation reduces toxic and pro-apoptotic effects of the amyloid beta-protein fragment (31–35) on isolated brain mitochondria, *Neuroscience* 126 (2004) 297–303.
- [147] J. Kanski, S. Varadarajan, M. Aksenova, D.A. Butterfield, Role of glycine-33 and methionine-35 in Alzheimer's amyloid beta-peptide 1–42-associated oxidative stress and neurotoxicity, *Biochim. Biophys. Acta* 1586 (2002) 190–198.
- [148] D. Pogocki, C. Schoneich, Redox properties of Met(35) in neurotoxic beta-amyloid peptide. A molecular modelling study, *Chem. Res. Toxicol.* 15 (2002) 408–418.
- [149] J. Naslund, A. Schierhorn, U. Hellman, L. Lannfelt, A.D. Roses, L.O. Tjebberg, J. Silberring, S.E. Gandy, B. Winblad, P. Greengard, et al., Relative abundance of Alzheimer A beta amyloid peptide variants in Alzheimer disease and normal aging, *Proc. Natl. Acad. Sci. U. S. A.* 91 (1994) 8378–8382.
- [150] Y.M. Kuo, T.A. Kokjohn, T.G. Beach, L.I. Sue, D. Brune, J.C. Lopez, W.M. Kalback, D. Abramowski, C. Sturchler-Pierrat, M. Staufenbiel, A.E. Roher, Comparative analysis of amyloid-beta chemical structure and

- amyloid plaque morphology of transgenic mouse and Alzheimer's disease brains, *J. Biol. Chem.* 276 (2001) 12991–12998.
- [151] V.K. Singh, J.F. Cheng, S.J. Leu, Effect of substance P and protein kinase inhibitors on beta-amyloid peptide-induced proliferation of cultured brain cells, *Brain Res.* 660 (1994) 353–356.
- [152] D.T. Stephenson, J.A. Clemens, In vivo effects of beta-amyloid implants in rodents: lack of potentiation of damage associated with transient global forebrain ischemia, *Brain Res.* 586 (1992) 235–246.
- [153] T. Takenouchi, E. Munekata, beta-Amyloid peptide, substance P, and SEC receptor ligand activate cytoplasmic Ca²⁺ in neutrophil-like HL-60 cells: effect of chemotactic peptide antagonist BocMLF, *Peptides* 16 (1995) 1019–1024.
- [154] J.S. Whitson, C.G. Glabe, E. Shintani, A. Abcar, C.W. Cotman, Beta-amyloid protein promotes neuritic branching in hippocampal cultures, *Neurosci. Lett.* 110 (1990) 319–324.
- [155] J.S. Whitson, D.J. Selkoe, C.W. Cotman, Amyloid beta protein enhances the survival of hippocampal neurons in vitro, *Science* 243 (1989) 1488–1490.
- [156] B.A. Yankner, L.K. Duffy, D.A. Kirschner, Neurotrophic and neurotoxic effects of amyloid beta protein: reversal by tachykinin neuropeptides, *Science* 250 (1990) 279–282.
- [157] M.A. Simmons, C.R. Schneider, Amyloid beta peptides act directly on single neurons, *Neurosci. Lett.* 150 (1993) 133–136.
- [158] K. Ueda, Y. Fukui, H. Kageyama, Amyloid beta protein-induced neuronal cell death: neurotoxic properties of aggregated amyloid beta protein, *Brain Res.* 639 (1994) 240–244.
- [159] M.P. Lambert, A.K. Barlow, B.A. Chromy, C. Edwards, R. Freed, M. Liosatos, T.E. Morgan, I. Rozovsky, B. Trommer, K.L. Viola, P. Wals, C. Zhang, C.E. Finch, G.A. Krafft, W.L. Klein, Diffusible, nonfibrillar ligands derived from Abeta_{1–42} are potent central nervous system neurotoxins, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 6448–6453.
- [160] C.J. Pike, A.J. Walencewicz, C.G. Glabe, C.W. Cotman, Aggregation-related toxicity of synthetic beta-amyloid protein in hippocampal cultures, *Eur. J. Pharmacol.* 207 (1991) 367–368.
- [161] C. Behl, J.B. Davis, F.G. Klier, D. Schubert, Amyloid beta peptide induces necrosis rather than apoptosis, *Brain Res.* 645 (1994) 253–264.
- [162] H.G. Lee, R.J. Castellani, X. Zhu, G. Perry, M.A. Smith, Amyloid-beta in Alzheimer's disease: the horse or the cart? Pathogenic or protective? *Int. J. Exp. Pathol.* 86 (2005) 133–138.
- [163] A. Kontush, C. Berndt, W. Weber, V. Akopyan, S. Arlt, S. Schipling, U. Beisiegel, Amyloid-beta is an antioxidant for lipoproteins in cerebrospinal fluid and plasma, *Free Radic. Biol. Med.* 30 (2001) 119–128.
- [164] C.-W. Chan, A. Dharmarajan, C.S. Atwood, X. Huang, R.E. Tanzi, A.I. Bush, R.N. Martins, Anti-apoptotic action of Alzheimer Abeta, *Alzheimer's Rep.* 2 (1999) 1–6.
- [165] J. Wang, D.W. Dickson, J.Q. Trojanowski, V.M. Lee, The levels of soluble versus insoluble brain Abeta distinguish Alzheimer's disease from normal and pathologic aging, *Exp. Neurol.* 158 (1999) 328–337.
- [166] T. Koppal, R. Subramaniam, J. Drake, M.R. Prasad, H. Dhillon, D.A. Butterfield, Vitamin E protects against Alzheimer's amyloid peptide (25–35)-induced changes in neocortical synaptosomal membrane lipid structure and composition, *Brain Res.* 786 (1998) 270–273.
- [167] W. Vogt, Oxidation of methionyl residues in proteins: tools, targets, and reversal, *Free Radic. Biol. Med.* 18 (1995) 93–105.
- [168] V. Kadlcik, C. Sicard-Roselli, T.A. Mattioli, M. Kodicek, C. Houee-Levin, One-electron oxidation of beta-amyloid peptide: sequence modulation of reactivity, *Free Radic. Biol. Med.* 37 (2004) 881–891.
- [169] S. Varadarajan, S. Yatin, J. Kanski, F. Jahanshahi, D.A. Butterfield, Methionine residue 35 is important in amyloid beta-peptide-associated free radical oxidative stress, *Brain Res. Bull.* 50 (1999) 133–141.
- [170] C.C. Page, C.C. Moser, X. Chen, P.L. Dutton, Natural engineering principles of electron tunnelling in biological oxidation–reduction, *Nature* 402 (1999) 47–52.
- [171] A. Rauk, D.A. Armstrong, D.P. Fairlie, Is oxidative damage mediated by amyloid beta and prion peptide mediated by hydrogen atom transfer from glycine alpha-carbon to methionine sulfur within beta-sheets? *J. Am. Chem. Soc.* 122 (2000) 9761–9767.
- [172] M.A. Lovell, W.D. Ehmann, M.P. Mattson, W.R. Markesbery, Elevated 4-hydroxynonenal in ventricular fluid in Alzheimer's disease, *Neurobiol. Aging* 18 (1997) 457–461.
- [173] H. Esterbauer, R.J. Schaur, H. Zollner, Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes, *Free Radic. Biol. Med.* 11 (1991) 81–128.
- [174] G. Spateller, Peroxyl radicals: inductors of neurodegenerative and other inflammatory diseases. Their origin and how they transform cholesterol, phospholipids, plasmalogens, polyunsaturated fatty acids, sugars, and proteins into deleterious products, *Free Radic. Biol. Med.* 41 (2006) 362–387.
- [175] M. Palmblad, A. Westlind-Danielsson, J. Bergquist, Oxidation of methionine 35 attenuates formation of amyloid beta -peptide 1–40 oligomers, *J. Biol. Chem.* 277 (2002) 19506–19510.
- [176] F.E. Ali, K.J. Barnham, C.J. Barrow, F. Separovic, Metal catalyzed oxidation of tyrosine residues by different oxidation systems of copper/hydrogen peroxide, *J. Inorg. Biochem.* 98 (2004) 173–184.
- [177] C.S. Atwood, X. Huang, A. Khatri, R.C. Scarpa, Y.S. Kim, R.D. Moir, R.E. Tanzi, A.E. Roher, A.I. Bush, Copper catalyzed oxidation of Alzheimer Abeta, *Cell Mol. Biol. (Noisy-le-grand)* 46 (2000) 777–783.
- [178] J.M. Onorato, S.R. Thorpe, J.W. Baynes, Immunohistochemical and ELISA assays for biomarkers of oxidative stress in aging and disease, *Ann. N. Y. Acad. Sci.* 854 (1998) 277–290.
- [179] C.S. Atwood, G. Perry, H. Zeng, Y. Kato, W.D. Jones, K.Q. Ling, X. Huang, R.D. Moir, D. Wang, L.M. Sayre, M.A. Smith, S.G. Chen, A.I. Bush, Copper mediates dityrosine cross-linking of Alzheimer's amyloid-beta, *Biochemistry* 43 (2004) 560–568.
- [180] D. Metodiewa, Molecular mechanisms of cellular injury produced by neurotoxic amino acids that generate reactive oxygen species, *Amino Acids* 14 (1998) 181–187.
- [181] S. Pennathur, V. Jackson-Lewis, S. Przedborski, J.W. Heinecke, Mass spectrometric quantification of 3-nitrotyrosine, ortho-tyrosine, and o,o'-dityrosine in brain tissue of 1-methyl-4-phenyl-1,2,3, 6-tetrahydropyridine-treated mice, a model of oxidative stress in Parkinson's disease, *J. Biol. Chem.* 274 (1999) 34621–34628.
- [182] D.A. Malencik, S.R. Anderson, Dityrosine as a product of oxidative stress and fluorescent probe, *Amino Acids* 25 (2003) 233–247.
- [183] Y.M. Kuo, M.R. Emmerling, C. Vigo-Pelfrey, T.C. Kasunic, J.B. Kirkpatrick, G.H. Murdoch, M.J. Ball, A.E. Roher, Water-soluble Abeta (N-40, N-42) oligomers in normal and Alzheimer disease brains, *J. Biol. Chem.* 271 (1996) 4077–4081.
- [184] A.E. Roher, M.O. Chaney, Y.M. Kuo, S.D. Webster, W.B. Stine, L.J. Haverkamp, A.S. Woods, R.J. Cotter, J.M. Tuohy, G.A. Krafft, B.S. Bonnell, M.R. Emmerling, Morphology and toxicity of Abeta-(1–42) dimer derived from neuritic and vascular amyloid deposits of Alzheimer's disease, *J. Biol. Chem.* 271 (1996) 20631–20635.
- [185] D.M. Walsh, I. Klyubin, J.V. Fadeeva, W.K. Cullen, R. Anwyl, M.S. Wolfe, M.J. Rowan, D.J. Selkoe, Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo, *Nature* 416 (2002) 535–539.
- [186] I. Yoon, K.H. Lee, J. Cho, Gossypin protects primary cultured rat cortical cells from oxidative stress- and beta-amyloid-induced toxicity, *Arch. Pharm. Res.* 27 (2004) 454–459.
- [187] P. Zatta, G. Tognon, P. Carampin, Melatonin prevents free radical formation due to the interaction between beta-amyloid peptides and metal ions [Al(III), Zn(II), Cu(II), Mn(II), Fe(II)], *J. Pineal. Res.* 35 (2003) 98–103.
- [188] Z. Feng, Y. Chang, Y. Cheng, B.L. Zhang, Z.W. Qu, C. Qin, J.T. Zhang, Melatonin alleviates behavioral deficits associated with apoptosis and cholinergic system dysfunction in the APP 695 transgenic mouse model of Alzheimer's disease, *J. Pineal. Res.* 37 (2004) 129–136.
- [189] E. Matsubara, T. Bryant-Thomas, J. Pacheco Quinto, T.L. Henry, B. Poeeggeler, D. Herbert, F. Cruz-Sanchez, Y.J. Chyan, M.A. Smith, G. Perry, M. Shoji, K. Abe, A. Leone, I. Grundke-Ikbal, G.L. Wilson, J. Ghiso, C. Williams, L.M. Refolo, M.A. Pappolla, D.G. Chain, E. Neria, Melatonin increases survival and inhibits oxidative and amyloid pathology in a transgenic model of Alzheimer's disease, *J. Neurochem.* 85 (2003) 1101–1108.
- [190] Y. Cheng, Z. Feng, Q.Z. Zhang, J.T. Zhang, Beneficial effects of

- melatonin in experimental models of Alzheimer disease, *Acta Pharmacol. Sin.* 27 (2006) 129–139.
- [191] F.V. Defeudis, Bilobalide and neuroprotection, *Pharmacol. Res.* 46 (2002) 565–568.
- [192] B.S. Oken, D.M. Storzbach, J.A. Kaye, The efficacy of Ginkgo biloba on cognitive function in Alzheimer disease, *Arch. Neurol.* 55 (1998) 1409–1415.
- [193] P.L. Le Bars, Response patterns of EGb 761 in Alzheimer's disease: influence of neuropsychological profiles, *Pharmacopsychiatry* 36 (Suppl. 1) (2003) S50–S55.
- [194] L.S. Schneider, S.T. DeKosky, M.R. Farlow, P.N. Tariot, R. Hoerr, M. Kieser, A randomized, double-blind, placebo-controlled trial of two doses of Ginkgo biloba extract in dementia of the Alzheimer's type, *Curr. Alzheimer Res.* 2 (2005) 541–551.
- [195] Z. Yao, Drieu.V. Papadopoulos, The Ginkgo biloba extract EGb 761 rescues the PC12 neuronal cells from beta-amyloid-induced cell death by inhibiting the formation of beta-amyloid-derived diffusible neurotoxic ligands, *Brain Res.* 889 (2001) 181–190.
- [196] G.J. Kelloff, J.A. Crowell, V.E. Steele, R.A. Lubet, W.A. Malone, C.W. Boone, L. Kopelovich, E.T. Hawk, R. Lieberman, J.A. Lawrence, I. Ali, J.L. Viner, C.C. Sigman, Progress in cancer chemoprevention: development of diet-derived chemopreventive agents, *J. Nutr.* 130 (2000) 467S–471S.
- [197] M. Ganguli, V. Chandra, M.I. Kamboh, J.M. Johnston, H.H. Dodge, B.K. Thelma, R.C. Juyal, R. Pandav, S.H. Belle, S.T. DeKosky, Apolipoprotein E polymorphism and Alzheimer disease: The Indo-US Cross-National Dementia Study, *Arch. Neurol.* 57 (2000) 824–830.
- [198] S. Shishodia, G. Sethi, B.B. Aggarwal, Curcumin: getting back to the roots, *Ann. N. Y. Acad. Sci.* 1056 (2005) 206–217.
- [199] M.Y. Aksenov, W.R. Markesbery, Changes in thiol content and expression of glutathione redox system genes in the hippocampus and cerebellum in Alzheimer's disease, *Neurosci. Lett.* 302 (2001) 141–145.
- [200] J.M. Ringman, S.A. Frautschy, G.M. Cole, D.L. Masterman, J.L. Cummings, A potential role of the curry spice curcumin in Alzheimer's disease, *Curr. Alzheimer Res.* 2 (2005) 131–136.
- [201] F. Yang, G.P. Lim, A.N. Begum, O.J. Ubeda, M.R. Simmons, S.S. Ambegaokar, P.P. Chen, R. Kaye, C.G. Glabe, S.A. Frautschy, G.M. Cole, Curcumin inhibits formation of amyloid beta oligomers and fibrils, binds plaques, and reduces amyloid in vivo, *J. Biol. Chem.* 280 (2005) 5892–5901.
- [202] L. Baum, A. Ng, Curcumin interaction with copper and iron suggests one possible mechanism of action in Alzheimer's disease animal models, *J. Alzheimers Dis.* 6 (2004) 367–377 (discussion 443–369).
- [203] H. Keberle, The biochemistry of desferrioxamine and its relation to iron metabolism, *Ann. N. Y. Acad. Sci.* 119 (1964) 758–768.
- [204] M.P. Cuajungco, K.Y. Faget, X. Huang, R.E. Tanzi, A.I. Bush, Metal chelation as a potential therapy for Alzheimer's disease, *Ann. N. Y. Acad. Sci.* 920 (2000) 292–304.
- [205] J.R. Walton, Aluminum in hippocampal neurons from humans with Alzheimer's disease, *Neurotoxicology* 27 (2006) 385–394.
- [206] E.K. Perry, R.H. Perry, New insights into the nature of senile (Alzheimer-type) plaques, *Trends Neurosci.* 8 (1985) 301–303.
- [207] D.P. Perl, A.R. Brody, Alzheimer's disease: X-ray spectrometric evidence of aluminum accumulation in neurofibrillary tangle-bearing neurons, *Science* 208 (1980) 297–299.
- [208] M.A. Lovell, W.D. Ehmann, W.R. Markesbery, Laser microprobe analysis of brain aluminum in Alzheimer's disease, *Ann. Neurol.* 33 (1993) 36–42.
- [209] M. Kawahara, Effects of aluminum on the nervous system and its possible link with neurodegenerative diseases, *J. Alzheimers Dis.* 8 (2005) 171–182 (discussion 209–115).
- [210] V. Moret, Y. Laras, N. Pietrancosta, C. Garino, G. Quelever, A. Rolland, B. Mallet, J.C. Norreel, J.L. Kraus, 1,1'-Xylyl bis-1,4,8,11-tetraaza cyclotetradecane: a new potential copper chelator agent for neuroprotection in Alzheimer's disease. Its comparative effects with clioquinol on rat brain copper distribution, *Bioorg. Med. Chem. Lett.* 16 (2006) 3298–3301.
- [211] J.Y. Lee, J.E. Friedman, I. Angel, A. Kozak, J.Y. Koh, The lipophilic metal chelator DP-109 reduces amyloid pathology in brains of human beta-amyloid precursor protein transgenic mice, *Neurobiol. Aging* 25 (2004) 1315–1321.
- [212] R.A. Cherny, C.S. Atwood, M.E. Xilinas, D.N. Gray, W.D. Jones, C.A. McLean, K.J. Barnham, I. Volitakis, F.W. Fraser, Y. Kim, X. Huang, L.E. Goldstein, R.D. Moir, J.T. Lim, K. Beyreuther, H. Zheng, R.E. Tanzi, C.L. Masters, A.I. Bush, Treatment with a copper-zinc chelator markedly and rapidly inhibits beta-amyloid accumulation in Alzheimer's disease transgenic mice, *Neuron* 30 (2001) 665–676.
- [213] M. Perluigi, G. Joshi, R. Sultana, V. Calabrese, C. De Marco, R. Coccia, D.A. Butterfield, In vivo protection by the xanthate tricyclodecan-9-yl-xanthogenate against amyloid beta-peptide (1–42)-induced oxidative stress, *Neuroscience* 138 (2006) 1161–1170.
- [214] S. Nakae, T. Iwa, Y. Asai, H. Anjiki, Y. Kameta, [Long-term follow-up following anastomosis of the superior vena cava and pulmonary artery], *Nippon Kyobu Geka Gakkai Zasshi* 21 (1973) 530–537.
- [215] M.S. Yassin, J. Ekblom, M. Xilinas, C.G. Gottfries, L. Oreland, Changes in uptake of vitamin B(12) and trace metals in brains of mice treated with clioquinol, *J. Neurol. Sci.* 173 (2000) 40–44.
- [216] C.W. Ritchie, A.I. Bush, A. Mackinnon, S. Macfarlane, M. Mastwyk, L. MacGregor, L. Kiers, R. Cherny, Q.X. Li, A. Tammer, D. Carrington, C. Mavros, I. Volitakis, M. Xilinas, D. Ames, S. Davis, K. Beyreuther, R.E. Tanzi, C.L. Masters, Metal-protein attenuation with iodochlorhydroxyquin (clioquinol) targeting Abeta amyloid deposition and toxicity in Alzheimer disease: a pilot phase 2 clinical trial, *Arch. Neurol.* 60 (2003) 1685–1691.
- [217] C. Reinhard, S.S. Hebert, B. De Strooper, The amyloid-beta precursor protein: integrating structure with biological function, *EMBO J.* 24 (2005) 3996–4006.